

**Table 1 Contribution to breeding of different breeding sites as reported by more than 500 support workers (“mosquito men”) throughout the year in Khartoum State, 2001–04**

<b>Year/breeding sites</b>	<b>Broken pipes</b>	<b>Indoor breeding</b>	<b>Construction basin</b>	<b>Farms/ irrigation</b>	<b>Institutes</b>	<b>Total</b>
<i>2001</i>						
No. sites inspected	215 814	322 911	489 814	365 641	460 676	1 854 856
No. with anopheline larvae	6 005	918	1 884	25 336	1 587	35 760
Contribution to breeding (%)	16.8	2.6	5.3	70.9	4.4	1.4
<i>2002</i>						
No. sites inspected	459 822	537 835	323 836	3 709 658	2 056 479	7 123 630
No. with anopheline larvae	4 501	431	920	75 263	1 969	83 084
Contribution to breeding (%)	5.4	0.5	1.1	90.6	2.4	1.2
<i>2003</i>						
No. sites inspected	666 403	727 110	241 628	5 646 925	1 454 820	8 736 886
No. with anopheline larvae	13 908	379	985	96 059	2 202	113 533
Contribution to breeding (%)	12.2	0.3	0.9	84.6	1.9	1.3
<i>2004</i>						
No. sites inspected	816 283	1 277 126	166 030	7 919 977	2 180 868	12 360 284
No. with anopheline larvae	7 739	388	210	148 888	1 406	158 631
Contribution to breeding (%)	4.9	0.2	0.1	93.9	0.9	1.2