

Table 3 Bacterial isolation according to characteristics of the patients with the diarrhoea episode

| Characteristic | Bacterial culture result | | Adjusted odds ratio | 95% CI | | |
|--------------------------------|--------------------------|------------|---------------------|--------|--------------|------------|
| | Negative No. | Negative % | | | Positive No. | Positive % |
| <i>Age group (years)</i> | | | | | | |
| < 1 | 445 | 94.7 | 25 | 5.3 | 0.2 | 0.08–0.50 |
| 1–3 | 145 | 75.9 | 46 | 24.1 | 1.3 | 0.5–3.2 |
| 3–5 | 84 | 62.7 | 50 | 37.3 | 2.4 | 0.9–6.1 |
| > 5 ^a | 47 | 77.0 | 14 | 23.0 | | |
| <i>Sex</i> | | | | | | |
| Male | 396 | 87.2 | 58 | 12.8 | 0.5 | 0.3–1.0 |
| Female ^a | 330 | 82.1 | 72 | 17.9 | | |
| <i>Level of dehydration</i> | | | | | | |
| Mild to moderate | 706 | 84.9 | 126 | 15.1 | 1.4 | 0.3–8.3 |
| Severe ^a | 20 | 83.3 | 4 | 16.7 | | |
| <i>Number of stools/day</i> | | | | | | |
| 1–5 | 278 | 84.8 | 50 | 15.2 | 0.7 | 0.3–1.6 |
| 6–9 | 311 | 86.1 | 50 | 13.9 | 0.7 | 0.3–1.6 |
| 10–30 ^a | 136 | 81.4 | 31 | 18.6 | | |
| <i>Associated illness</i> | | | | | | |
| Yes | 74 | 77.1 | 22 | 22.9 | 1.8 | 0.8–4.2 |
| No ^a | 652 | 85.8 | 108 | 14.2 | | |
| <i>Prior antibiotics given</i> | | | | | | |
| No | 650 | 85.5 | 110 | 14.5 | 0.4 | 0.2–1.1 |
| Yes ^a | 76 | 79.2 | 20 | 20.8 | | |
| <i>Blood in stool</i> | | | | | | |
| Yes | 98 | 65.3 | 52 | 34.7 | 3.0 | 1.6–5.7 |
| No ^a | 628 | 89.0 | 78 | 11.0 | | |

^aReference group.

CI = confidence interval.