

Table 2 Prevalence of tuberculosis (TB) among household TB contacts screened for active TB according to age group and category of TB, Morocco 1993–2004

Year	Any form of TB (%)				SSPPT (%)				SSNPT (%)				EPT (%)				SPC (%)			
	Age (years)			P	Age (years)			P	Age (years)			P	Age (years)			P	Age (years)			P
	All	< 10	≥ 10		All	< 10	≥ 10		All	< 10	≥ 10		All	< 10	≥ 10		All	<10	≥10	
1993	2.0	1.4	2.3	<0.001	1.0	0.1	1.3	<0.001	0.4	0.1	0.5	<0.001	0.3	0.1	0.4	<0.001	0.4	1.1	0.1	<0.001
1994	3.1	2.0	3.5	<0.01	1.8	0.1	2.3	<0.001	0.3	0.1	0.3	<0.001	0.5	0.1	0.6	<0.001	0.5	1.7	0.2	<0.001
1995	2.1	2.1	2.1	NS	1.0	0.1	1.2	<0.001	0.3	0.1	0.4	<0.001	0.3	0.2	0.3	<0.001	0.5	1.7	0.1	<0.001
1996	2.4	1.7	2.7	<0.001	1.0	0.1	1.4	<0.001	0.4	0.1	0.5	<0.001	0.5	0.2	0.6	<0.001	0.5	1.4	0.3	<0.001
1997	3.7	2.2	4.3	<0.001	1.7	0.1	2.3	<0.001	0.5	0.1	0.6	<0.001	0.9	0.3	1.2	<0.001	0.6	1.7	0.2	<0.001
1998	1.9	1.8	1.9	NS	0.8	0.0	1.0	<0.001	0.2	0.1	0.2	<0.001	0.5	0.2	0.6	<0.001	0.4	1.5	0.1	<0.001
1999	1.9	1.9	1.9	NS	0.8	0.0	1.0	<0.001	0.3	0.1	0.4	<0.001	0.3	0.1	0.4	<0.001	0.4	1.6	0.1	<0.001
2000	2.3	2.9	2.2	<0.001	0.9	0.1	1.1	<0.001	0.3	0.1	0.3	<0.001	0.6	0.4	0.6	<0.001	0.5	2.4	0.1	<0.001
2001	3.1	2.1	3.3	<0.001	1.5	0.1	1.7	<0.001	0.4	0.2	0.4	<0.001	0.9	0.4	1.0	<0.001	0.4	1.4	0.2	<0.001
2002	2.4	2.5	2.4	NS	1.1	0.2	1.3	<0.001	0.3	0.2	0.3	NS	0.6	0.5	0.7	<0.05	0.4	1.6	0.1	<0.001
2003	2.2	1.8	2.3	<0.01	1.1	0.2	1.3	<0.001	0.2	0.1	0.3	<0.001	0.5	0.4	0.6	<0.01	0.3	1.2	0.1	<0.001
2004	2.0	1.9	2.0	NS	1.0	0.1	1.1	<0.001	0.2	0.1	0.2	NS	0.5	0.3	0.6	<0.01	0.3	1.4	0.1	<0.001
Overall	2.5	2.0	2.6	<0.001	1.1	0.1	1.5	<0.001	0.3	0.1	0.4	<0.001	0.5	0.2	0.6	<0.001	0.5	1.6	0.1	<0.001

P-value assesses statistical significance of the proportion difference between the age groups < 10 and ≥ 10.

NS = not statistically significant (P ≥ 0.05).

SSPPT = sputum smear-positive pulmonary TB; SSNPT: sputum smear-negative pulmonary TB; EPT = extrapulmonary TB; SPC = symptomatic primary complex.