

Table 4 Clients' motives and sources of information about voluntary counselling and testing (VCT) services by clients' sex

Variable	Males (n = 759)		Females (n = 169)		χ^2 -value	P-value
	No.	%	No.	%		
<i>Motives for using the service</i>						
Risky sexual behaviour	298	39.3	19	11.2	48.25	0.001
Previous blood transfusion	169	22.3	38	22.5	0.004	0.951
Planning to get married	120	15.8	23	13.6	0.51	0.474
Injecting drug user	129	17.0	2	1.2	28.51	0.001
New sexual partner	122	16.1	7	4.1	16.44	0.001
Partner has risky sexual behaviour	56	7.4	57	33.7	89.74	0.001
Referred by health worker	65	8.6	15	8.9	0.02	0.896
Previous HIV testing	29	3.8	8	4.7	0.30	0.583
Other	47	6.2	43	25.4	58.50	0.001
<i>Source of information about VCT</i>						
Relative/friend	243	32.0	60	35.5	0.76	0.382
Poster	183	24.1	44	26.0	0.28	0.599
Health worker	172	22.7	45	26.6	1.21	0.271
Lecture	155	20.4	31	18.3	0.37	0.542
Another VCT client	163	21.5	17	10.1	11.52	0.001
Pamphlet	130	17.1	20	11.8	2.86	0.091
Sex partner/spouse	52	6.9	13	7.7	0.15	0.698
Television	47	6.2	9	5.3	0.18	0.669
Newspaper	15	2.0	4	2.4	0.11	0.746
Telephone hotline	16	2.1	3	1.8	0.08	0.782
Radio	13	1.7	5	3.0	1.13	0.288
Other	139	18.3	19	11.2	4.89	0.027