

Table 2 Antibiotic resistance patterns of *Mycobacterium tuberculosis* genotypes by patient's sex and by region of the Syrian Arab Republic

Variable	Antibiotic resistance pattern ^a												
	RRRR	RRSS	RRSR	RRRS	SSRS	SRSS	SRRR	SRRS	RSSS	RSRS	SRSR	SSRR	SSSS
No. of isolates													
Male	24	5	5	6	8	2	5	2	2	1	0	0	3
Female	15	0	0	0	6	0	1	0	0	1	1	1	0
Total	39	5	5	6	14	2	6	2	2	2	1	1	3
Genotypes by region													
Aleppo	<i>G10, G25, G30, G44, G45, G47, G51, G52, G59, G61, G67, G71 G72, G74, G82, Gx</i>	<i>G20, G31</i>	<i>G43, G20</i>	-	<i>G3, G19, G23, G27 G28, G32</i>	<i>G21</i>	<i>G23; G49; G70; Gx</i>	<i>G54</i>	<i>G57</i>	-	<i>G20</i>	-	<i>G32</i>
Damascus	<i>G20, G23, G29, G53, G78, G81, Gx</i>	-	-	<i>G20, Gx</i>	<i>G20</i>	-	-	-	<i>G3</i>	<i>G55</i>	-	-	<i>G45</i>
Deir ez-Zor	<i>G20, G60, Gx</i>	-	<i>G20</i>	<i>G79</i>	-	-	<i>G3</i>	<i>G42</i>	-	-	-	<i>G35</i>	<i>G22</i>
Rif Dimashq	<i>G23</i>	-	-	-	<i>G76</i>	-	-	-	-	-	-	-	-
Latakia	<i>G3, G38</i>	-	-	<i>G62</i>	<i>Gx</i>	<i>G48</i>	-	-	-	-	-	-	-
Homs	<i>G21</i>	-	<i>G22</i>	-	<i>G80</i>	-	-	-	-	<i>G20</i>	-	-	-
Al-Hasakah	<i>G40</i>	<i>G24</i>	-	-	<i>G37</i>	-	-	-	-	-	-	-	-
Hama	<i>G20</i>	-	-	<i>G26</i>	<i>G43</i>	-	-	-	-	-	-	-	-
Al-Qamishli	<i>G36, G20</i>	-	-	-	-	-	-	-	-	-	-	-	-
Ar-Raqqah	<i>G50</i>	-	-	-	-	-	<i>G56</i>	-	-	-	-	-	-
Abu Kamal	-	<i>G24</i>	-	-	-	-	-	-	-	-	-	-	-
Adra	-	<i>G34</i>	-	-	-	-	-	-	-	-	-	-	-
As-Suwayda'	-	-	-	-	<i>G64</i>	-	-	-	-	-	-	-	-
Quneitra	<i>G33</i>	-	-	-	-	-	-	-	-	-	-	-	-
Iraq	<i>G69, G75, G77</i>	-	<i>G22</i>	<i>G65</i>	-	-	-	-	-	-	-	-	-

^aAntibiotic resistance pattern sequence: rifampin (RIF)-isoniazid (INH)-streptomycin (STR)-ethambutol (EMB).

R = resistant; S = sensitive.

G = genotype (bold = isolates from females); Gx = no genotype pattern found; - = no isolates.