

Table 2 Prevalence and factors associated with hepatitis B surface antigen (HBsAg) in Pakistan by age, marital status and history of injection use

Variable	Males		Females		Total	
	No. (%)	OR (95%CI)	No. (%)	OR (95% CI)	No.(%)	OR (95% CI)
Age (years)						
< 5	33 (1.6)	1	20 (1.0)	1	53 (1.3)	1
5-19	198 (2.1)	1.33 (0.90-1.96)	131 (1.5)	1.56 (0.95-2.57)	329 (1.8)	1.42 (1.05-1.92)
20-29	104 (2.4)	1.55 (1.03-2.35)	88 (2.1)	2.17 (1.30-3.64)	192 (2.2)	1.79 (1.30-2.46)
30-39	106 (3.7)	2.45 (1.63-3.72)	81 (2.8)	2.91 (1.74-4.91)	187 (3.2)	2.61 (1.90-3.60)
40-49	104 (4.5)	2.97 (1.97-4.50)	63 (3.0)	3.11 (1.83-5.34)	167 (3.8)	3.04 (2.20-4.20)
50-59	82 (5.5)	3.70 (2.41-5.68)	34 (2.6)	2.72 (1.51-4.92)	116 (4.1)	3.37 (2.40-4.75)
≥ 60	82 (5.0)	3.32 (2.16-5.10)	30 (2.7)	2.79 (1.53-5.13)	112 (4.0)	3.28 (2.33-4.62)
Total	709 (3.0)		447 (2.0)		1156 (2.5)	
Marital status (20-49 years)						
Never married	67 (1.9)	1	36 (1.7)	1	103 (1.8)	1
Married	245 (4.2)	2.32 (1.75-3.09)	193 (2.8)	1.70 (1.70-2.47)	438 (3.4)	1.96 (1.57-2.45)
Divorced/separated/widowed	2 (2.7)	1.49 (0.0-6.3)	3 (1.5)	0.88 (0.27-2.84)	5 (1.8)	1.01 (0.36-2.59)
Total	314 (3.3)		232 (2.5)		546 (2.9)	
No. of intramuscular injections						
None	111 (1.9)	1	87 (1.7)	1	198 (1.8)	1
< 5	359 (3.1)	1.68 (1.35-2.09)	215 (2.0)	1.13 (0.87-1.47)	574 (2.5)	1.42 (1.20-1.68)
5-10	177 (3.3)	1.83 (1.43-2.34)	114 (2.2)	1.87 (1.54-2.26)	291 (2.8)	1.55 (1.29-1.81)
> 10	62 (4.4)	2.43 (1.75-3.37)	31 (2.0)	1.17 (0.76-1.80)	93 (3.2)	1.78 (1.38-2.30)
Total	709 (2.9)		447 (2.0)		1156 (2.5)	
Types of syringes						
None	115 (1.9)	1	83 (1.7)	1	198 (1.8)	1
New disposable	201 (2.5)	1.35 (1.06-1.71)	124 (1.6)	0.96 (0.72-1.29)	325 (2.1)	1.16 (0.97-1.39)
Re-used syringes	332 (3.7)	1.99 (1.60-2.49)	206 (2.4)	1.40 (1.08-1.83)	538 (3.0)	1.71 (1.44-2.02)
Don't know	65 (4.6)	2.48 (1.80-3.42)	30 (2.2)	1.30 (0.83-2.02)	95 (3.4)	1.92 (1.48-2.47)
Total	706 (2.9)		450 (2.0)		1156 (2.5)	

CI = confidence interval; OR = odds ratio.