

Table 1 Distribution of various dermatoses by age and sex of the studied patients (n = 633) (concluded)

| Type of dermatosis | 1-3 years | | | | 4-7 years | | | | 8-12 years | | | | All ages | |
|---|-----------|-----|-------|-----|-----------|-----|-------|-----|------------|-----|-------|-----|----------|------|
| | M | F | Total | | M | F | Total | | M | F | Total | | | |
| | No. | No. | No. | % | No. | No. | No. | % | No. | No. | No. | % | No. | % |
| Pigmentary (17.8%) | | | | | | | | | | | | | | |
| Vitiligo | 0 | 3 | 3 | 0.4 | 10 | 12 | 22 | 3.3 | 15 | 15 | 30 | 4.6 | 55 | 8.3 |
| Post-inflammatory hypopigmentation | 7 | 4 | 11 | 1.7 | 10 | 10 | 20 | 3.0 | 6 | 0 | 6 | 0.9 | 37 | 5.6 |
| Post-inflammatory hyperpigmentation | 1 | 5 | 6 | 0.9 | 6 | 6 | 12 | 1.8 | 8 | 0 | 8 | 1.2 | 26 | 3.9 |
| Total | 8 | 12 | 20 | 3.0 | 26 | 28 | 54 | 8.1 | 29 | 15 | 44 | 6.7 | 118 | 17.8 |
| Papulosquamous (14.2%) | | | | | | | | | | | | | | |
| Psoriasis | 5 | 0 | 5 | 0.8 | 7 | 3 | 10 | 1.5 | 10 | 10 | 20 | 3.0 | 35 | 5.3 |
| Lichen planus | 3 | 0 | 3 | 0.4 | 2 | 10 | 12 | 1.8 | 6 | 5 | 11 | 1.7 | 26 | 3.9 |
| Pityriasis rosea | 2 | 2 | 4 | 0.6 | 6 | 4 | 10 | 1.5 | 0 | 5 | 5 | 0.8 | 19 | 2.9 |
| Pityriasis rubra | 0 | 0 | 0 | - | 4 | 0 | 4 | 0.6 | 6 | 4 | 10 | 1.5 | 14 | 2.1 |
| Pilaris | | | | | | | | | | | | | | |
| Total | 10 | 2 | 12 | 1.8 | 19 | 17 | 36 | 5.4 | 22 | 24 | 46 | 7.0 | 94 | 14.2 |
| Drug-induced (4.5%) | | | | | | | | | | | | | | |
| Total | 5 | 2 | 7 | 1.1 | 10 | 3 | 13 | 1.9 | 7 | 3 | 10 | 1.5 | 30 | 4.5 |
| Nutritional deficiency (1.8%) | | | | | | | | | | | | | | |
| Hair involvement (Flag sign, easily pluckable, sparse and depigmented hair) | 6 | 2 | 8 | 1.2 | 0 | 0 | 0 | - | 0 | 0 | 0 | - | 8 | 1.2 |
| Skin involvement (flaky and hypopigmented skin) | 4 | 0 | 4 | 0.6 | 0 | 0 | 0 | - | 0 | 0 | 0 | - | 4 | 0.6 |
| Total | 10 | 2 | 12 | 1.8 | 0 | 0 | 0 | - | 0 | 0 | 0 | - | 12 | 1.8 |

M = males; F = females.