30-39 11 16 ≥40 9 9 Woman's education Illiterate

those who did not (n = 34)

Woman's age (years)

Variable

19-29

1-12 years > 12 years

Residency Urban

Rural

< 352

≥ 353

Gravidity

1-3

4-14

Yes

No

Yes

No

Yes

No

≤10

> 10

Yes

No

2-37

38-43

Yes

No

No

symptoms Yes

Easy access to hospital

Gestational age (weeks)

Refused termination of pregnancy

Unrecognized pregnancy danger

Family size Mean (SD)

Pregnancy planned

Health insurance

Prenatal care in current pregnancy

Distance to nearest hospital by car (min)

5 1 11

15

Yes

17

Table 2 Demographic, pregnancy and access variables of maternal deaths in women who delayed seeking care (n = 42) and

No

14

9

25 9

12

11

13

17

17

3

22

8

5.1(2.3)

19

15

6

22

0

12

22

1

33

1

33

OR (95% CI)

NA

NA

1.42 (0.96-2.09)

1.36 (0.78-2.38)

0.99 (0.63-1.56)

0.79 (0.50-1.24)

1.10 (0.64-1.91)

0.65 (0.46-0.91)

0.60 (0.37-0.98)

0.50 (0.37-0.67)

1.78 (1.12-2.82)

1.85 (1.36-2.52)

6.32 (3.02-13.2)

P-value^b

0.828

0.303

0.092

0.254

0.969

0.360

0.722

< 0.001

0.078

0.035

0.006

0.008

0.010

< 0.001

Delay in seeking care

8

23

Family income (US\$ monthly)

19 19 9

15 20

5.6 (2.9)

26

8

12

17

22

9

27

14

10

32

36

6

"Numbers may not add to 76 due to missing data; "Chi-squared test for binomial and t-test for means. OR = unadjusted odds ratio; CI = confidence interval.; NA = not applicable; SD = standard deviation.

23

8 27 8