

Table 1 Sociodemographic characteristics and outcome of cases of suicide (attempted and completed)

Variable	All cases		Attempted (nonfatal) (n = 49 217)		Completed (fatal) (n = 3883)		Case fatality rate ^a	Completed suicide OR (95% CI)	P-value
	No.	%	No.	%	No.	%			
Total	53 100	100.0	49 217	100.0	3 883	100.0	7.3		
Sex									
Female	29 225	55.0	27 312	55.5	1 913	49.3	6.5	0.79 (0.74–0.84)	< 0.001
Male	23 652	44.5	21 759	44.2	1 893	48.8	8.0	Reference	
Missing data	223	0.4	146	0.3	77	2.0	–		
Age (years)									
≤ 20	19 832	37.3	18 659	37.9	1 173	30.2	5.9	0.43 (0.37–0.49)	< 0.001
21–30	21 148	39.8	19 753	40.1	1 395	35.9	6.6	0.45 (0.39–0.51)	< 0.001
31–40	5 415	10.2	4 909	10.0	506	13.0	9.3	0.63 (0.55–0.74)	< 0.001
41–50	2 740	5.2	2 425	4.9	315	8.1	11.5	0.81 (0.68–0.96)	< 0.001
≥ 51	2 477	4.7	2 145	4.4	332	8.6	13.4	Reference	
Missing data	1 488	2.8	1 326	2.7	162	4.2	–	–	
Marital status									
Single	16 499	31.1	15 216	30.9	1 283	33.1	7.8	0.39 (0.22–0.68)	< 0.001
Married	12 165	22.9	10 793	21.9	1 372	35.4	11.3	0.16 (0.09–0.28)	< 0.001
Divorced	71	0.1	54	0.1	17	0.4	23.9	Reference	
Missing data	24 365	45.9	23 154	47.1	1 211	31.1	–	–	
Suicide method									
Drug overdose	38 784	73.0	37 880	77.0	904	23.3	2.3	0.01 (0.01–0.01)	< 0.001
Cutting	763	1.4	734	1.5	29	0.7	3.8	0.01 (0.01–0.04)	< 0.05
Poisoning	5 626	10.6	5 531	10.9	275	7.1	4.9	0.02 (0.01–0.03)	< 0.001
Narcotic overdose	359	0.7	319	0.6	40	1.0	11.1	0.06 (0.04–0.10)	< 0.001
Self-burning	2 164	4.1	713	1.4	1 451	37.4	67.1	0.73 (0.49–1.08)	< 0.05
Hanging	1 239	2.3	291	0.6	948	24.4	76.5	1.16 (0.77–1.75)	< 0.05
Firearm	130	0.2	34	0.1	96	2.5	73.8	Reference	
Other methods	441	0.8	375	0.8	66	1.7	14.9	–	
Missing data	3 594	6.8	3 520	7.2	74	1.9	–	–	

^aCase fatality rate = [fatal/fatal + nonfatal] × 100.

OR = odds ratio; CI = confidence interval.