

**Table 4 Distribution of body mass index (BMI) categories by scores on the total Eating Attitudes Test (EAT-26) scale and subscales of the Eating Disorder Inventory-3 (EDI-3) in male adolescents (2003 and 2010 cohorts combined)**

Scale	Underweight		Normal		Overweight		Obese		$\chi^2$
	No.	%	No.	%	No.	%	No.	%	
<b>Total EAT-26</b>									
Normal score	34	77.3	450	84.4	122	76.3	34	66.7	13.72*
High score (> 20)	10	22.7	83	15.6	38	23.8	17	33.3	
<b>Drive for thinness</b>									
Normal score	45	97.8	583	95.4	168	84.4	52	77.6	45.52***
High score ( $\geq 17$ )	1	2.2	28	4.6	31	15.6	15	22.4	
<b>Bulimia</b>									
Normal score	40	83.3	468	74.4	137	66.5	38	58.5	13.87***
High score ( $\geq 5$ )	8	16.7	161	25.6	69	33.5	27	41.5	
<b>Body dissatisfaction</b>									
Normal score	46	95.8	581	97.2	155	79.1	33	51.6	166.2***
High score ( $\geq 22$ )	2	4.2	17	2.8	41	20.9	31	48.4	
<b>Low self-esteem</b>									
Normal score	42	87.5	534	85.9	178	85.2	52	78.8	2.57
High score ( $\geq 9$ )	6	12.5	88	14.1	31	14.8	14	21.2	

Chi-squared tests: \* $P < 0.05$ ; \*\*\* $P < 0.001$ .