

**Table 6 Distribution of studied female sex workers in relation to their knowledge about male condoms**

Variable	No.	%
<i>Places where condoms can be obtained (n = 424)</i>		
Pharmacies	315	74.3
Supermarkets	1	0.2
Clinics	4	0.9
Hospitals	14	3.3
Bars/hotels	3	0.7
From a friend	15	3.5
Nongovernmental organizations	23	5.4
Don't know	100	23.6
<i>Have used a condom more than once (n = 429)</i>		
Yes	9	2.1
No	420	97.9
<i>Have used a lubricant with condoms (n = 427)</i>		
Yes	40	9.4
No	387	90.6
<i>Have experienced condom tear (n = 429)</i>		
Yes	55	12.8
No	374	87.2
<i>Response to condom tear (n = 55)</i>		
Stopped sexual act	5	9.1
Continued after using a new condom	34	61.8
Continued without condom	16	29.1
<i>Had sex during menses (n = 431)</i>		
Yes	128	29.7
No	303	70.3
<i>Used condom for sex during menses (n = 128)</i>		
Yes	45	35.1
No	79	61.7
No answer	4	3.1
<i>Group sex<sup>a</sup></i>		
Have had group sex (n = 431)	33	7.7
Used condom during group sex (n = 33)	7	21.2

<sup>a</sup>Having sex with more than 1 partner at the same time.