

Table 1 Species and sites of the adult anopheline mosquitoes in Bashagard district by collection method

Species	Village	Collection method					No.	%
		Pit shelter	Hand catch	Total catch	Night-biting catch	Larval rearing		
<i>Anopheles culicifacies</i>	Tisur	4	5	24	19	16	68	9.81
	Daranar	7	11	25	26	31	100	14.43
<i>An. dthali</i>	Tisur	2	8	73	13	42	138	19.92
	Daranar	3	6	82	31	29	151	21.79
<i>An. fluviatilis</i>	Tisur	-	-	-	8	-	8	1.16
	Daranar	-	-	-	3	-	3	0.43
<i>An. moghulensis</i>	Tisur	-	-	-	4	31	35	5.05
	Daranar	-	-	-	3	6	9	1.30
<i>An. stephensi</i>	Tisur	6	7	42	2	11	68	9.81
	Daranar	3	5	21	4	15	48	6.93
<i>An. superpictus</i>	Tisur	5	4	-	6	3	18	2.59
	Daranar	4	3	1	3	15	26	3.75
<i>An. turkhudi</i>	Tisur	-	-	-	10	2	12	1.73
	Daranar	-	2	-	3	4	9	1.30
Total		34	51	268	135	205	693	100