

Table 3 History of lid surgery for trachomatous trichiasis (TT) of participants' eyes in different demographic groups, by blindness status (n = 8191 eyes)

Variable	Blind (vision < 3/60)		Not blind (vision ≥ 3/60)		OR (95% CI)
	No.	%	No.	%	
Total group					
Surgery history -ve	600	97.4	7217	95.3	0.54 (0.32-0.89)
Surgery history +ve	16	2.6	358	4.7	
Total	616	100.0	7575	100.0	
Males					
Surgery history -ve	342	98.8	2628	97.1	0.39 (0.12-0.99)
Surgery history +ve	4	1.2	78	2.9	
Subtotal	346	100.0	2706	100.0	
Females					
Surgery history -ve	258	95.6	4589	94.2	0.76 (0.40-1.33)
Surgery history +ve	12	4.4	280	5.8	
Subtotal	270	100.0	4869	100.0	
Hyperendemic area					
Surgery history -ve	248	93.9	6413	94.8	1.03 (0.60-1.80)
Surgery history +ve	14	5.3	350	5.2	
Subtotal	364	100.0	6661	100.0	
Mesoendemic area					
Surgery history -ve	352	99.4	804	99.0	0.57 (0.12-2.70)
Surgery history +ve	2	0.6	8	1.0	
Subtotal	354	100.0	812	100.0	

Data presented are numbers and % of eyes examined.
OR = odds ratio; CI = confidence interval.