

Table 4 Likelihood of calling smoking cessation clinics and hotlines among current smokers in relation to sociodemographic characteristics

Variable	Likelihood of calling a smoking cessation clinic or hotline							
	Very likely		Likely		Unlikely		Very unlikely	
	No.	%	No.	%	No.	%	No.	%
Total	132	12.8	415	40.2	249	24.1	236	22.9
Sex								
Male	113	12.5	370	40.9	219	24.3	202	22.4
Female	19	15.0	45	35.4	30	22.8	34	26.8
				$\chi^2 = 2.37; P = 0.499$				
Age (years)								
18–29	47	15.9	119	40.3	81	27.5	48	16.3
30–39	53	15.8	140	41.7	62	18.5	81	24.1
40–49	26	10.4	93	37.2	67	26.8	64	25.6
50–59	2	2.4	38	45.8	18	21.7	25	30.1
60+	4	5.9	25	36.8	21	30.9	18	26.5
				$\chi^2 = 34.40; P < 0.001 (r = 0.13; P < 0.001)$				
Education								
Elementary or less	17	10.2	73	43.7	36	21.6	41	24.6
Preparatory to high school	83	13.5	261	42.6	137	22.3	132	21.5
Diploma and above	32	12.2	81	30.9	76	29.0	73	27.9
				$\chi^2 = 16.64; P = 0.011 (r = 0.06; P = 0.076)$				
Monthly income (JD)^a								
< 300	56	11.2	208	41.5	140	27.9	97	19.4
300–599	71	18.2	149	38.2	74	19.0	96	24.6
600+	5	3.7	53	39.3	35	25.9	42	31.1
				$\chi^2 = 33.94; P < 0.001 (r = 0.07; P = 0.038)$				
Ever tried to quit smoking								
No	31	23.3	117	28.3	122	49.2	115	48.5
Yes, but failed	102	76.7	297	71.7	126	50.8	122	51.5
				$\chi^2 = 53.36; P < 0.001 (r = -0.21; P < 0.001)$				

^a6 respondents refused to declare their income.