

Table 4 Antimicrobial susceptibility of multidrug resistant *Acinetobacter baumannii* isolates (n = 116) to various antibiotics tested

Variable	No. of isolates	Susceptibility (%) ^c
Multidrug resistant ^a	90	77.6
Colistin + minocycline	37	31.9
Colistin + tobramycin	1	0.9
Colistin + sulfamethoxazole/trimethoprim	12	10.3
Colistin + ampicillin/sulbactam	6	5.2
Colistin + ampicillin/sulbactam + sulfamethoxazole/trimethoprim	2	1.7
Colistin + sulfamethoxazole/trimethoprim + tobramycin	1	0.9
Colistin + minocycline + ampicillin/sulbactam	1	0.9
Colistin + minocycline + sulfamethoxazole/trimethoprim	9	7.8
Colistin + minocycline + tetracycline	10	8.6
Colistin + minocycline + tobramycin	2	1.7
Colistin + minocycline + tetracycline + tobramycin + sulfamethoxazole/trimethoprim	9	7.8
Pandrug resistant ^b	10	8.6
Not susceptible to any of the tested antibiotics	2	1.7
Susceptible only to colistin	8	6.9

^aResistant to imipenem plus 3 or more different antibiotic classes.; ^bResistant to all tested antibiotics or only susceptible to colistin; ^cPercentage of total number of isolates tested (n = 116) susceptible to colistin only.