

Table 1 Annual isolation rate of non-O1 and non-O139 *Vibrio cholerae* at Aga Khan University clinical laboratory; 1999-2012

Year of culture testing	Stool culture performed	Stool culture growing diarrhoeal pathogens	<i>V. cholerae</i> among diarrhoeal pathogens	<i>V. cholerae</i> strains				
				O1 (Ogawa)	O1 (Inaba)	O139	Non-O1 and non-O139	%
	No.	No.	No.	No.	No.	No.	No.	%
1999	4 807	841	317	282	31	0	4	1
2000	4 806	735	271	213	1	39	18	7
2001	5 587	1108	412	325	2	81	4	1
2002	8 062	1 334	229	142	3	55	29	13
2003	9 943	2 063	262	182	9	2	69	27
2004	4 700	851	98	64	19	1	14	14
2005	5 556	2 473	738	2	721	0	15	2
2006	6 103	1 441	188	0	181	0	7	4
2007	6 365	1 084	286	149	122	0	15	5
2008	8 267	1 884	238	219	3	0	16	7
2009	9 999	2 667	248	223	5	0	20	8
2010	7 224	1 343	155	143	3	0	9	6
2011	7 597	1 294	152	136	10	0	6	4
2012	6 784	1 007	74	15	52	0	7	10
Total	95 800	20 124	3668	2095	1162	178	233	6