

Table 4 Distribution of the studied women according to admission data and outcome

Admission data	Survived (<i>n</i> = 410)		Died (<i>n</i> = 38)		Total (<i>n</i> = 448)		<i>P</i> -value
	No.	%	No.	%	No.	%	
<i>Admission diagnosis</i>							< 0.001 ²
Severe pre-eclampsia/eclampsia	176	96.2	7	3.8	183	40.8	
Severe postpartum haemorrhage	100	96.2	4	3.8	104	23.2	
Severe systemic infection or sepsis	4	22.2	14	77.8	18	4.0	
Ruptured uterus	9	81.8	2	18.2	11	2.5	
Other diagnoses ¹	121	91.7	11	8.3	132	29.5	
<i>Length of stay in ICU (days)</i>							< 0.001 ³
Mean (SD)		2.56 (1.7)		8.71 (8.21)		3.07 (3.33)	
Range		1–14		1–34		1–34	
<i>Pregnancy state on admission</i>							0.034 ⁴
Pregnant	181	94.8	10	5.2	191	42.6	
Delivered	229	89.1	28	10.9	257	57.4	

¹Other diagnoses include heart diseases with pregnancy, acute fatty liver of pregnancy, respiratory complications, antepartum haemorrhage, ectopic pregnancy, complications of abortion, vesicular mole, systemic lupus erythematosus, diabetic ketoacidosis and deep vein thrombosis.

²Chi-squared = 120; ³Mann-Whitney *Z* = -4.61; ⁴Monte Carlo test.

ICU = intensive care unit, SD = standard deviation.