

Table 1 Anthropometric and clinical data for patients with suspected or confirmed diagnosis of small molecule disorders on admission to a paediatric intensive care unit (PICU) (*n* = 34)

Characteristic	Value
Age at PICU presentation (months)	
Min-max	1.17-72.00
Median (IQR)	9.00 (2.88-24.00)
Sex, No. (%)	
Males	15 (44.1 %)
Females	19 (55.9 %)
Residence, No. (%)	
Rural	19 (55.9 %)
Urban	15 (44.1 %)
Weight (kg)	
Min-max	1.55-20
Median (IQR)	7.00 (4.78-10.25)
Paediatric Index of Mortality 2 (on admission)	
Min-max	20-90.80
Mean (SD)	53.10 (27.14)
Median (IQR)	44.95 (27.60-85.00)
Length of stay (d)	
Min-max	1-26
Mean (SD)	6.97 (6.11)
Median (IQR)	3.50 (2.00-10.00)
PELOD score on day 2^a	
Min-max	4-31
Mean (SD)	13.52 (8.03)
Median (IQR)	10.00 (5.00-20.00)
Outcome, No. (%)	
Discharged	16 (47.1)
Died	11 (32.3)
Died ≤ 24 hours following admission	7 (20.6)

IQR = interquartile range.

PELOD score = Paediatric Logistic Organ Dysfunction score.

SD = standard deviation.

^a7 cases died ≤ 24 hrs of admission

Min = minimum; Max = maximum