

Table 3 Subgroups of anthropometric failure according to CIAF

Group	Description	Age (mo)						Total age and sex combined n = 1292 (100%)
		Boys		Total	Girls		Total	
		6-23 n = 138	24-59 n = 543	6-59 n = 681	6-23 n = 141	24-59 n = 470	6-59 n = 611	
A	No failure	48 (34.8)	158 (29.0)	206 (30.2)	64 (45.4)	117 (24.9)	181 (29.6)	387 (29.9)
B	Wasting only	11 (07.9)	58 (10.7)	69 (10.1)	06 (04.3)	59 (12.6)	65 (10.6)	134 (10.4)
C	Wasting/ underweight	30 (21.7)	77 (14.2)	107 (15.7)	13 (09.2)	77 (16.4)	90 (14.7)	197 (15.3)
D	Wasting/stunting/ underweight	15 (10.9)	84 (15.5)	99 (14.5)	12 (08.5)	74 (15.7)	86 (14.1)	185 (14.3)
E	Stunting/ underweight	15 (10.9)	115 (21.2)	130 (19.1)	28 (19.9)	95 (20.2)	123 (20.1)	253 (19.6)
F	Stunting only	09 (06.5)	26 (04.8)	35 (05.1)	08 (05.7)	16 (03.4)	24 (03.9)	59 (4.6)
Y	Underweight only	10 (07.2)	25 (04.6)	35 (05.1)	10 (07.0)	32 (06.8)	42 (06.9)	77 (5.9)
Total CIAF (B+C+D+E+F+Y)		90 (65.2)	385 (70.9)	475 (69.8)	77 (54.6)	353 (75.1)	430 (70.4)	905 (70.1)
Single failure (B+F+Y)								270 (20.9)
Multiple failures (C+E+D)								635 (49.2)

Values in parentheses are percentages.