

Table 2 Comparison of responses on selected variables between 2009 and 2015 by EMR Member Dates

Selected variables	Result in 2015	Result in 2009*
FP safety regulation at Health facility α	16/16 (100%)	16/18 (89%)
FP guidelines and protocols updated β	15/16 (94%)	13/18 (72%)
FP as basic package and delivered at Primary Health Care level	16/16 (100%)	17/18 (94%)
Mix contraceptives in essential drug list	16/16 (100%)	16/18 (89%)
FP counselling and methods in preconception care	6/16 (38%)	NA
FP counselling by nurse	14/16 (88%)	14/18 (78%)
FP effective supply chain management all over the country	13/16 (81%)	12/18 (67%)
FP pre- and in-service training programme in universities	14/15 (93%)	14/18 (78%)
Training guidelines and materials: evidence-based	14/15 (93%)	15/18 (83%)
FP special program for adolescents	8/16 (50%)	6/18 (33%)
FP special program for displaced or refugees	12/13 (92%)	9/18 (50%)
FP special program for poor/disadvantaged	10/15 (67%)	9/18 (50%)
FP promoted through Effective social marketing	10/15 (67%)	12/18 (67%)
FP promoted through community mobilization efforts	12/15 (80%)	11/18 (61%)
HMIS system collects and analyses FP data	15/16 (94%)	13/18 (72%)

*Please note the 2009 data are taken from a previous regional survey on FP (8).

α : This refers to utilization of WHO MEC standards in prescription of contraceptives to clients.