

Table 1 Usage of antibiotics by indication

Indication* N (%)	PPS1	PPS2	PPS3	PPS4	PPS5	PPS6
HEENT infection	2 (5.1%)	2 (5.4%)	2 (5.9%)	2 (5.3%)	2 (5.3%)	0 (0.0%)
SSS infection	1 (2.6%)	2 (5.4%)	4 (12.0%)	6 (16.0%)	6 (16.0%)	5 (15.0%)
CAP	1 (2.6%)	3 (8.1%)	1 (2.9%)	2 (5.3%)	2 (5.3%)	0 (0.0%)
HAP	2 (5.1%)	2 (5.4%)	1 (2.9%)	1 (2.6%)	1 (2.6%)	2 (6.1%)
PJP prophylaxis	3 (7.7%)	3 (8.1%)	4 (12.0%)	3 (7.9%)	3 (7.9%)	3 (9.1%)
GI infection	6 (15.0%)	5 (14.0%)	6 (18.0%)	6 (16.0%)	3 (7.9%)	3 (9.1%)
UT infection (complicated/uncomplicated cystitis/ catheter-acquired)	2 (5.1%)	2 (5.4%)	1 (2.9%)	1 (2.6%)	1 (2.6%)	2 (6.1%)
Blood infection (bacteremia)	8 (21.0%)	6 (16.0%)	5 (15.0%)	5 (13.0%)	4 (11.0%)	4 (12.0%)
Febrile neutropenia	8 (21.0%)	7 (19.0%)	8 (24.0%)	8 (21.0%)	7 (18.0%)	7 (21.0%)
Undefined/unknown	6 (15.0%)	5 (14.0%)	2 (5.9%)	4 (11.0%)	9 (24.0%)	7 (21.0%)
Total	39	37	34	38	38	33

*HEENT: head-eye-ear-nose-throat; SSS: skin and skin structure; CAP: community-acquired pneumonia; HAP: hospital-acquired pneumonia; PJP: *Pneumocystis jirovecii* Pneumonia; GI: gastrointestinal; UT: urinary tract