

Table 2 Association of current midwakh use with other variables related to tobacco uses: logistic regression analyses

Variable	Midwakh use	OR (95% CI)	P-value	ORa (95% CI)a	P-value
	No. (%)				
<i>Tried a cigarette before age 14 years (of those who ever tried a cigarette)</i>					
No	39 (7.5)	1 (-)		1 (-)	
Yes	138 (15.7)	2.31 (1.53-3.49)	0.001*	2.34 (1.49-3.68)	0.001*
<i>Current cigarette smoker</i>					
No	90 (1.6)	1 (-)		1 (-)	
Yes	161 (21.6)	16.61 (12.21-22.58)	< 0.001*	15.22 (11.08-20.90)	< 0.001*
<i>Waterpipe use</i>					
Never	34 (1)	1 (-)		1 (-)	
Ever	228 (9)	9.89 (6.78-14.45)	< 0.001*	9.61 (6.66-13.86)	< 0.001*
<i>Parent/guardian smokes (any type)</i>					
No	86 (3.3)	1 (-)		1 (-)	
Yes	168 (5.2)	1.61 (1.13-2.29)	0.01*	1.56 (1.05-2.31)	0.029*
<i>Exposure to second-hand smoke</i>					
No	59 (3.6)	1 (-)		1 (-)	
Yes	206 (4.8)	1.37 (0.94-2.0)	0.097	1.32 (0.91-1.92)	0.128

OR: unadjusted odds ratio; ORa: adjusted odds ratio; CI confidence intervals.

*Controlling for age, sex and school type.

*Statistically significant at P < 0.05.