

Table 3 Prevalence of underweight and overweight by age and sex according to international reference cut-offs

Variable	Prevalence (%)												P-value
	Girls						Boys						
	6 years (n = 46)	7 years (n = 69)	8 years (n = 44)	9 years (n = 42)	10 years (n = 53)	Total (n = 254)	6 years (n = 65)	7 years (n = 48)	8 years (n = 57)	9 years (n = 41)	10 years (n = 44)	Total (n = 255)	
Underweight													
WHO	0.0	0.0	0.0	2.4	3.8	1.1	0.0	0.0	3.5	2.4	2.3	1.6	0.71
IOTF	8.7	13.0	4.5	9.5	15.1	10.6	9.2	8.3	5.3	7.3	4.5	7.1	0.15
CDC	4.3	2.9	2.3	2.4	9.4	4.3	1.5	2.1	3.5	2.4	2.3	2.4	0.21
French reference	4.4	1.5	0.0	2.4	3.8	2.4	0.0	2.1	3.5	2.4	2.3	2.0	0.75
Overweight													
WHO	23.9	18.8*	34.1	23.8	22.6*	24.0*	32.3	39.6	22.8	26.8	43.2	32.6	0.03
IOTF	21.7	18.8	29.5	19.0	18.9	21.3	24.6	27.1	19.3	21.9	29.5	24.3	0.41
CDC	21.7	17.4*	27.3	16.7	15.1	19.3	24.6	37.5	19.3	21.9	29.5	26.3	0.06
French reference	17.4	14.5	27.3	19.1	18.9	18.9	23.1	22.9	14.0	21.9	29.5	22.0	0.39

WHO: World Health Organization, IOTF: International Obesity Task Force, CDC: Centers for Disease Control and Prevention.

*Significant at $P < 0.05$.