

Table 4 Cancer deaths attributable to inadequate physical activity by age, sex and cancer site, Tunisia, 2019

Age, in years	Post-menopausal breast			Lung			Colon/rectum			Endometrium			Stomach			Kidney			Bladder			Oesophagus		
	AD	PAF	OD	AD	PAF	OD	AD	PAF	OD	AD	PAF	OD	AD	PAF	OD	AD	PAF	OD	AD	PAF	OD	AD	PAF	OD
Men																								
35-44	NA	NA	NA	9	16.7	55	2	NE	17	NA	NA	NA	2	NE	13	1	NE	3	1	NE	4	1	NE	2
45-64	NA	NA	NA	155	17.0	909	20	11.9	166	NA	NA	NA	20	17.0	119	4	11.3	36	14	18.0	80	6	19.9	33
≥ 65	NA	NA	NA	224	17.3	1293	46	12.1	378	NA	NA	NA	45	17.4	257	7	11.6	64	61	18.5	329	11	20.3	56
Total	NA	NA	NA	388	17.2	2257	68	12.0	561	NA	NA	NA	67	17.3	389	12	11.4	103	76	18.4	413	18	20.1	91
Women																								
35-44	NA	NA	NA	0	NE	12	3	11.9	26	0	NE	2	2	NE	14	0	NE	2	0	NE	1	0	NE	1
45-64	50	12.9	386	15	16.6	88	22	12.3	175	5	NE	24	11	17.6	64	2	NE	13	2	NE	12	2	NE	8
≥ 65	51	12.9	395	26	16.9	155	48	12.6	378	9	19.9	45	26	17.9	148	3	11.9	25	10	18.9	53	4	NE	19
Total	101	12.9	781	43	16.8	255	73	12.5	579	14	19.8	71	39	17.8	226	5	11.9	40	12	18.9	66	6	20.8	28
Both																								
35-44	NA	NA	NA	11	16.6	67	5	11.7	43	0	NE	2	4	17.0	27	1	NE	5	1	NE	5	1	NE	3
45-64	50	12.9	386	169	17.0	997	42	12.1	341	5	NE	24	31	17.2	183	6	11.4	49	17	18.1	92	8	20.0	41
≥ 65	51	12.9	395	250	17.3	1448	94	12.4	756	9	19.9	45	71	17.6	405	10	11.7	89	71	18.5	382	15	20.5	75
Total	101	12.9	781	431	17.2	2512	141	12.3	1140	14	19.8	71	106	17.5	615	17	11.6	143	88	18.4	479	24	20.3	119

AD= attributable deaths. PAF= population attributable fraction. OD= observed deaths; NA= not applicable; NE= not estimated because the total (denominator) was less than 25 so no percentages could be calculated.

Note: AD and OD are numbers; PAF is a percentage.