

Table 3. Pregnancy and antenatal visits during the COVID-19 pandemic (March 2020–March 2021) (Ankara, 2021)

Pregnancy (n = 616)	n	%
<i>No pregnancy</i>	397	64.4
<i>Pregnancy</i>	219	35.6
Currently pregnant	86	14.0
Not currently pregnant	133	21.6
Livebirth	104	16.9
Abortion	29	4.7
Spontaneous	25	4.1
Induced	4	0.6
Characteristics	n	%
<i>Antenatal care (n = 219)</i>		
Yes	129	58.9
No	90	41.1
<i>Number of antenatal visits^a (n = 124^b)</i>		
Less than recommended number	50	40.3
Recommended number or more	74	59.7
<i>Place of antenatal care (n=129)</i>		
Migration health centre	91	70.5
Public hospital	48	37.2
Private practice/private hospital	12	9.3
House	3	2.3
<i>Antenatal care provider (n = 129)</i>		
Syrian doctor	92	71.3
Turkish physician	54	41.9
Turkish nurse	12	9.3
Syrian nurse	11	8.5
Syrian midwife	4	3.1
Turkish midwife	1	0.8
<i>Components of antenatal care^{c,d}</i>		
Iron pill supplementation (n = 129)	126	97.7
Blood pressure measuring (n = 126)	121	96.0
Blood test (n = 128)	121	95.3
Urinalysis (n = 128)	121	95.3
Measuring weight (n = 126)	112	88.9
Listening to foetal heartbeat (n = 126)	103	84.1
Nutritional recommendations (n = 129)	80	62.0

^aRecommended number in Türkiye ≥ 4 . No. of antenatal visits: mean 4.5 (3.0); median 4; 1st–3rd quartiles 3–6; range 1–25

^bNo response for 5 participants.

^cPercentage evaluated at least once. ^dDifferent numbers of unanswered questions.