

Table1 Distribution of respondents as regards awareness, use and support of condoms by sex, residence and schooling

Variable	Sex (n = 2534)			Residence (general population respondents only) (n = 1607)			Ever attended school (n = 2534)		
	Males %	Females %	P-value	Urban %	Rural %	P-value	Yes %	No %	P-value
<i>Awareness</i>									
Ever heard of HIV/AIDS	94.2	80.5	< 0.0001	90.5	81.0	< 0.0001	93.8	72.5	< 0.0001
Ever heard of the male condom	55.6	41.0	< 0.0001	55.0	45.3	< 0.0001	55.8	31.3	< 0.0001
Heard of the male condom and know it can prevent STIs	44.0	41.3	0.358	39.8	45.2	0.135	44.2	37.1	0.045
Mentioned male condom among known family planning/birth spacing methods	22.9	14.1	< 0.0001	23.1	18.4	0.028	23.3	7.6	< 0.0001
<i>Use in preceding 12 months</i>									
Used a method of family planning/ birth spacing in previous 12 months	23.7	35.7	< 0.0001	30.8	28.9	0.276	30.4	29.1	0.347
Used a male condom for family planning/birth spacing in previous 12 months	13.4	9.6	0.256	14.0	7.8	0.156	11.9	8.3	0.225
Used male condom for any reason during previous 12 months (married respondents only (n = 1586))	6.0	6.3	0.913	8.2	3.6	0.004	7.8	3.5	< 0.0001
<i>Support promotion of condom use</i>									
Supported condom promotion for those persisting with extramarital sex	24.2	23.0	< 0.0001	23.4	17.3	0.009	25.5	19.4	< 0.0001

STI = sexually transmitted infection.