

Table 6 Site-specific oral health conditions among adolescent smokeless tobacco users in Jeddah

Variable	Sextant ^c					
	A	B	C	D	E	F
Tobacco dipping site ^{a,b} (%)	5	95	5*	8*	30**	6***
Mean plaque score	1.81	1.47	1.88	1.53	1.45	1.53
Mean pocket depth (mm)	2.24	1.78	2.23	2.06	1.74	2.03
Mean clinical attachment loss (mm)	1.25	0.87	1.24	1.05	0.89	1.02
Mean gingival recession (mm)	0.01*	0	0.01*	0	0.03	0
Bleeding on probing (%)	47*	33	44	37	39	41
Mean DMFT score	0.34	0.39	0.39	0.47	0.04	0.46
D teeth (%)	83	93	86	70	100	80*
M teeth (%)	9	2	5	20	0	10
F teeth (%)	8	5*	9*	10*	0	10**
Oral mucosal lesions ^b (%)	2	96	4	2	12	2

^aSite of use of smokeless tobacco.

^bTotals more than 100% because some students use more than 1 site for dipping and have more than 1 lesion in the same sextant.

^cSextants A (from upper right second molar to upper right first premolar), B (from upper right cuspid to upper left cuspid), C (from upper left first premolar to upper left second molar), D (from lower left second molar to lower left first premolar), E (from lower left cuspid to lower right cuspid) and F (from lower left first premolar to lower left second molar).

*P < 0.05; **P < 0.01; ***P < 0.001.

D = decayed, M = missing, F = filled.