

Monitor children nutritional status, early detection of malnutrition and case referral. Share information and trigger alert verification..

Methodology

51%

Underwight

6

Exclusive

Breastfeeding

Nutritional assessment in NSS is normally performed through the measurement of child's weigh, height/length, MUAC, age, hemoglobin level, breastfeeding status and oedema. It is Facility-based surveillance system, established mainly in district. Additional screening of 21,581 children via mobile teams and pediatric wards integrated with TFC to ensure assessment of hospitals across the country.

children screened. It is not a formal survey using sampling methods to produce representative results.

The results only reflect the status of children who have attended the health facility in the month under review. The system tracks Global core nutrition indicators: Stunting, Wasting, Exclusive breastfeeding and Anemia. This bulletin is based on reports received by the health facility surveillance MoPHP teams*.

WHO supports the MoPHP and constantly works to expand coverage and improve data quality, funded by World Bank.

children in remote areas is conducted with ensuring to refer children with malnutrition forms to the nearest nutrition services to continue receiving treatment and follow up

Proportion of Severe Acute Malnutrition (SAM)

5% 5%

affected by SAM in June

< 115 mm

40%

Anemia Jun

- Moatsim Hael



Summary for Nutrition Indicators Jun 2024 in Reporting Gov

Acute Malnutrition by MUAC / Gender (Monthly Trend)

Proportion of Moderate Acute Malnutrition (MAM)


< 125 mm and >= 115 mm

40%


30%

affected by MAM in June

				Acute Malnutrition by Z- Score				Acute Malnutriton by MUAC				Underweight		Stunting		Exclusive Breastfeeding		Anemia		Oedema	
Governorate	# of Screened Children (0 - 5)	# of Screened Children (6 - 59)	Propartion of GAM (0 - 5)	# of MAM	Proportion of MAM	# of SAM	Proportion of SAM	# of MAM		# of SAM	Proportion of SAM	# of Underweight	Proportion of Under weight	# of Stunting	Proportion of Stunting		Proportion of EBF (Yes)	# of Anemia	Proportion of Anemia	# of Oedema	Proportion of Oedema
Al Bayda	707	2,660	15.6%	390	15%	176	7%	358	13%	115	4%	1,286	38%	953	36%	99	12%	296	16%	1	0.0%
Al Dhale'e	133	688	17.3%	153	22%	67	10%	143	21%	79	11%	404	49%	318	46%	39	27%	54	13%	1	0.1%
Al Hudaydah	539	3,938	21.5%	1,333	34%	793	20%	1,291	33%	720	18%	3,184	71%	2,191	56%	61	9%	1,541	41%	0	0.0%
Al Jawf	273	1,522	17.2%	274	18%	126	8%	210	14%	78	5%	805	44%	600	39%	60	19%	477	33%	2	0.1%
Al Mahwit	892	2,893	16.0%	520	18%	176	6%	390	13%	164	6%	1,677	44%	1,235	43%	189	19%	608	21%	3	0.1%
Amanat Al Asimah	1,745	4,685	17.9%	1,085	23%	283	6%	520	11%	181	4%	2,939	46%	2,066	44%	274	15%	326	12%	0	0.0%
Amran	1,015	4,030	17.2%	879	22%	339	8%	908	23%	396	10%	2,711	54%	2,164	54%	105	9%	1,013	25%	1	0.0%
Dhamar	968	3,697	22.6%	655	18%	368	10%	815	22%	415	11%	2,795	60%	2,212	60%	81	7%	518	14%	11	0.2%
Hadramaut	881	1,138	13.8%	165	14%	70	6%	105	9%	26	2%	509	25%	259	23%	184	20%	107	40%	0	0.0%
Hajjah	1,134	5,876	11.6%	1,004	17%	356	6%	974	17%	381	6%	3,569	51%	2,923	50%	113	8%	1,798	31%	0	0.0%
lbb	1,775	4,068	20.8%	988	24%	398	10%	712	18%	292	7%	3,003	51%	1,854	46%	262	13%	612	15%	28	0.5%
Lahj	165	799	17.6%	137	17%	74	9%	99	12%	49	6%	409	42%	213	27%	55	28%	154	21%	0	0.0%
Marib	123	1,075	17.9%	125	12%	54	5%	110	10%	34	3%	370	31%	330	31%	58	40%	88	11%	1	0.1%
Raymah	229	1,362	30.6%	267	20%	92	7%	255	19%	80	6%	1,092	69%	927	68%	2	1%	389	29%	0	0.0%
Sa'ada	463	3,568	18.6%	907	25%	385	11%	959	27%	448	13%	2,530	62%	2,030	57%	94	14%	479	18%	2	0.0%
Sana'a	676	3,114	19.7%	608	20%	216	7%	523	17%	167	5%	1,858	49%	1,423	46%	137	18%	477	16%	1	0.0%
Aden	313	525	27.8%	118	22%	56	11%	93	18%	31	6%	358	43%	178	34%	141	43%			0	0.0%
Taizz	1,398	3,295	17.9%	797	24%	288	9%	606	18%	207	6%	2,335	50%	1,575	48%	304	20%	465	16%	4	0.1%
Abyan	48	471	25.0%	134	28%	69	15%	67	14%	20	4%	300	58%	211	45%	19	34%	21	27%	0	0.0%
Shabwah	64	382	26.6%	60	16%	16	4%	40	10%	9	2%	154	35%	134	35%	-	-	223	59%	0	0.0%




Chronic Malnutrition (Stunting) by Height/Age Z Score / Gender (Monthly Trend)



16%

14%





© World Health Organization 2019

Proportion of Oedema per month